arXiv:1411.7299v1 [math.CA] 26 Nov 2014

Two-variable −1 Jacobi polynomials

Vincent X. Genest[1], Jean-Michel Lemay[1], Luc Vinet[1], and Alexei Zhedanov[2]

[1]Centre de recherches mathématiques, Université de Montréal, P.O. Box 6128, Centre-ville Station, Montréal, Canada, H3C 3J7
[2]Donetsk Institute for Physics and Technology, Donetsk 340114, Ukraine

Abstract

A two-variable generalization of the Big −1 Jacobi polynomials is introduced and characterized. These bivariate polynomials are constructed as a coupled product of two univariate Big −1 Jacobi polynomials. Their orthogonality measure is obtained. Their bispectral properties (eigenvalue equations and recurrence relations) are determined through a limiting process from the two-variable Big q-Jacobi polynomials of Lewanowicz and Woźny. An alternative derivation of the weight function using Pearson-type equations is presented.

Keywords: Bivariate Orthogonal Polynomials, Big −1 Jacobi polynomials
AMS classification numbers: 33C50

1 Introduction

The purpose of this paper is to introduce and study a family of bivariate Big −1 Jacobi polynomials. These two-variable polynomials, which shall be denoted by $\mathscr{J}_{n,k}(x,y)$, depend on four real parameters $\alpha,\beta,\gamma,\delta$ such that $\alpha,\beta,\gamma>-1$, $\delta\neq 1$ and are defined as

$$\mathscr{J}_{n,k}(x,y)=J_{n-k}\left(y;\alpha,2k+\beta+\gamma+1,(-1)^k\delta\right)\rho_k(y)J_k\left(\frac{x}{y};\gamma,\beta,\frac{\delta}{y}\right), \tag{1}$$

with $k=0,1,\ldots$ and $n=k,k+1,\ldots$, where

$$\rho_k(y)=\begin{cases} y^k\left(1-\frac{\delta^2}{y^2}\right)^{\frac{k}{2}}, & k \text{ even},\\ y^k\left(1-\frac{\delta^2}{y^2}\right)^{\frac{k-1}{2}}\left(1+\frac{\delta}{y}\right), & k \text{ odd},\end{cases}$$

and where $J_n(x;a,b,c)$ denotes the one-variable Big −1 Jacobi polynomials [20] (see section 1.1). It will be shown that these polynomials are orthogonal with respect to a positive measure defined on the disjoint union of four triangular domains in the real plane. The polynomials $\mathscr{J}_{n,k}(x,y)$ will also be identified as a $q\to -1$ limit of the two-variable Big q-Jacobi polynomials introduced by Lewanowicz and Woźny in [16], which generalize the bivariate little q-Jacobi polynomials introduced by Dunkl in [3]. The bispectral properties of the Big −1 Jacobi polynomials will be determined from this identification. The polynomials $\mathscr{J}_{n,k}(x,y)$ will be shown to satisfy an explicit vector-type three term recurrence relation and it will be seen that they are the joint eigenfunctions of a pair of commuting first order differential operators involving reflections. By solving the Pearson-type system of equations arising from the symmetrization of these differential/difference operators, the weight function for the polynomials $\mathscr{J}_{n,k}(x,y)$ will be recovered.

The defining formula of the two-variable Big -1 Jacobi polynomials (1) is reminiscent of the expressions found in [15] for the Krall-Sheffer polynomials [14], which, as shown in [11], are directly related to two-dimensional superintegrable systems on spaces with constants curvature (see [17] for a review of superintegrable systems). The polynomials $\mathscr{J}_{n,k}(x,y)$ do not belong to the Krall-Sheffer classification, as they will be seen to obey *first order* differential equations with reflections. The results of [11] however suggest that the polynomials $\mathscr{J}_{n,k}(x,y)$ could be related to two-dimensional integrable systems with reflections such as the ones recently considered in [5, 7, 6, 9]. This fact motivates our examination of the polynomials $\mathscr{J}_{n,k}(x,y)$.

1.1 The Big -1 Jacobi polynomials

Let us now review some of the properties of the Big -1 Jacobi polynomials which shall be needed in the following. The Big -1 Jacobi polynomials, denoted by $J_n(x;a,b,c)$, were introduced in [20] as a $q=-1$ limit of the Big q-Jacobi polynomials [12]. They are part of the Bannai-Ito scheme of -1 orthogonal polynomials [10, 8, 18, 19]. They are defined by

$$J_n(x;a,b,c)=\begin{cases} {}_2F_1\left(\begin{matrix}-\frac{n}{2},\frac{n+a+b+2}{2}\\ \frac{a+1}{2}\end{matrix}\,\middle|\,\frac{1-x^2}{1-c^2}\right)+\frac{n(1-x)}{(1+c)(a+1)}\,{}_2F_1\left(\begin{matrix}1-\frac{n}{2},\frac{n+a+b+2}{2}\\ \frac{a+3}{2}\end{matrix}\,\middle|\,\frac{1-x^2}{1-c^2}\right), & n \text{ even},\\ {}_2F_1\left(\begin{matrix}-\frac{n-1}{2},\frac{n+a+b+1}{2}\\ \frac{a+1}{2}\end{matrix}\,\middle|\,\frac{1-x^2}{1-c^2}\right)-\frac{(n+a+b+1)(1-x)}{(1+c)(a+1)}\,{}_2F_1\left(\begin{matrix}-\frac{n-1}{2},\frac{n+a+b+3}{2}\\ \frac{a+3}{2}\end{matrix}\,\middle|\,\frac{1-x^2}{1-c^2}\right), & n \text{ odd}, \end{cases} \tag{2}$$

where ${}_2F_1$ is the standard Gauss hypergeometric function [1]; when no confusion can arise, we shall simply write $J_n(x)$ instead of $J_n(x;a,b,c)$. The polynomials (2) satisfy the recurrence relation

$$x\,J_n(x)=A_n\,J_{n+1}(x)+(1-A_n-C_n)\,J_n(x)+C_n\,J_{n-1}(x),$$

with coefficients

$$A_n=\begin{cases}\frac{(n+a+1)(c+1)}{2n+a+b+2}, & n \text{ even},\\ \frac{(1-c)(n+a+b+1)}{2n+a+b+2}, & n \text{ odd},\end{cases}\qquad C_n=\begin{cases}\frac{n(1-c)}{2n+a+b}, & n \text{ even},\\ \frac{(n+b)(1+c)}{2n+a+b}, & n \text{ odd}.\end{cases}$$

It can be seen that for $a,b>-1$ and $|c|\neq 1$ the polynomials $J_n(x)$ are positive-definite. The Big -1 Jacobi polynomials satisfy the eigenvalue equation

$$\mathscr{L}J_n(x)=\left\{(-1)^n\,(n+a/2+b/2+1/2)\right\}J_n(x), \tag{3}$$

where $\mathscr{L}$ is the most general first-order differential operator with reflection preserving the space of polynomials of a given degree. This operator has the expression

$$\mathscr{L}=\left[\frac{(x+c)(x-1)}{x}\right]\partial_x R+\left[\frac{c}{2x^2}+\frac{ca-b}{2x}\right](R-\mathbb{I})+\left[\frac{a+b+1}{2}\right]R,$$

where R is the reflection operator, i.e. $Rf(x)=f(-x)$, and $\mathbb{I}$ stands for the identity. The orthogonality relation of the Big -1 Jacobi polynomials is as follows. For $|c|<1$, one has

$$\int_{\mathscr{C}} J_n(x;a,b,c)\,J_m(x;a,b,c)\,\omega(x;a,b,c)\,\mathrm{d}x=\left[\frac{(1-c^2)^{\frac{a+b+2}{2}}}{(1+c)}\right]h_n(a,b)\,\delta_{nm}, \tag{4}$$

where the interval is $\mathscr{C}=[-1,-|c|]\cup[|c|,1]$ and the weight function reads

$$\omega(x;a,b,c)=\theta(x)(1+x)(x-c)(x^2-c^2)^{\frac{b-1}{2}}\,(1-x^2)^{\frac{a-1}{2}}, \tag{5}$$

with $\theta(x)$ is the sign function. The normalization factor h_n is given by

$$h_n(a,b)=\begin{cases}\dfrac{2\,\Gamma\left(\frac{n+b+1}{2}\right)\Gamma\left(\frac{n+a+3}{2}\right)\left(\frac{n}{2}\right)!}{(n+a+1)\,\Gamma\left(\frac{n+a+b+2}{2}\right)\left(\frac{a+1}{2}\right)_{\frac{n}{2}}^2}, & n \text{ even},\\ \dfrac{(n+a+b+1)\,\Gamma\left(\frac{n+b+2}{2}\right)\Gamma\left(\frac{n+a+2}{2}\right)\left(\frac{n-1}{2}\right)!}{2\,\Gamma\left(\frac{n+a+b+3}{2}\right)\left(\frac{a+1}{2}\right)_{\frac{n+1}{2}}^2}, & n \text{ odd},\end{cases} \tag{6}$$

where $(a)_n$ stands for the Pochhammer symbol [1]. For $|c|>1$, one has

$$\int_{\widetilde{\mathscr{C}}} J_n(x;a,b,c)\,J_m(x;a,b,c)\,\widetilde{\omega}(x;a,b,c)\,\mathrm{d}x=\left[\frac{\theta(c)(c^2-1)^{\frac{a+b+2}{2}}}{1+c}\right]\widetilde{h}_n(a,b)\,\delta_{nm}, \tag{7}$$

where the interval is $\widetilde{\mathscr{C}}=[-|c|,-1]\cup[1,|c|]$ and the weight function reads

$$\widetilde{\omega}(x;a,b,c)=\theta(c\,x)(1+x)(c-x)(c^2-x^2)^{\frac{b-1}{2}}\,(x^2-1)^{\frac{a-1}{2}}. \tag{8}$$

In this case the normalization factor has the expression

$$\widetilde{h}_n(a,b)=\begin{cases}\dfrac{2\,\Gamma\left(\frac{n+b+1}{2}\right)\Gamma\left(\frac{n+a+3}{2}\right)\left(\frac{n}{2}\right)!}{(n+a+1)\Gamma\left(\frac{n+a+b+2}{2}\right)\left(\frac{a+1}{2}\right)_{\frac{n}{2}}^2}, & n \text{ even},\\ \dfrac{(n+a+b+1)\Gamma\left(\frac{n+b+2}{2}\right)\Gamma\left(\frac{n+a+2}{2}\right)\left(\frac{n-1}{2}\right)!}{2\,\Gamma\left(\frac{n+a+b+3}{2}\right)\left(\frac{a+1}{2}\right)_{\frac{n+1}{2}}^2} & n \text{ odd}.\end{cases} \tag{9}$$

The normalization factors h_n and $\widetilde{h}_n$ were not derived in [20]. They have been obtained here using the orthogonality relation for the Chihara polynomials provided in [8] and the fact that the Big -1 Jacobi polynomials are related to the latter by a Christoffel transformation. The details of this derivation are presented in appendix A.

2 Orthogonality of the two-variable Big -1 Jacobi polynomials

We now prove the orthogonality property of the two-variable Big -1 Jacobi polynomials.

Proposition 2.1. *Let $\alpha,\beta,\gamma>-1$ and $|\delta|<1$. The two-variable Big -1 Jacobi polynomials defined by* (1) *satisfy the orthogonality relation*

$$\int_{D_y}\int_{D_x} \mathscr{J}_{n,k}(x,y)\;\mathscr{J}_{m,\ell}(x,y)\,W(x,y)\,\mathrm{d}x\,\mathrm{d}y=H_{nk}\;\delta_{k\ell}\delta_{nm}, \tag{10}$$

with respect to the weight function

$$W(x,y)=\theta(x\,y)|y|^{\beta+\gamma}(1+y)\left(1+\frac{x}{y}\right)\left(\frac{x-\delta}{y}\right)(1-y^2)^{\frac{\alpha-1}{2}}\left(1-\frac{x^2}{y^2}\right)^{\frac{\gamma-1}{2}}\left(\frac{x^2-\delta^2}{y^2}\right)^{\frac{\beta-1}{2}}. \tag{11}$$

The integration domain is prescribed by

$$D_x=[-|y|,-|\delta|]\cup[|\delta|,|y|],\qquad D_y=[-1,-|\delta|]\cup[|\delta|,1], \tag{12}$$

and the normalization factor H_{nk} has the expression

$$H_{nk}=\left[\frac{(1-\delta^2)^{\frac{2k+\alpha+\beta+\gamma+3}{2}}}{(1+(-1)^k\delta)}\right]\,h_k(\gamma,\beta)\,h_{n-k}(\alpha,2k+\gamma+\beta+1),$$

where $h_n(a,b)$ is given by (6).

Proof. We proceed by a direct calculation. We denote the orthogonality integral by

$$I = \int\limits_{D_y}\int\limits_{D_x} \mathscr{J}_{n,k}(x,y)\, \mathscr{J}_{m,\ell}(x,y)\, W(x,y)\, \mathrm{d}x\, \mathrm{d}y.$$

Upon using the expressions (1) and (11) in the above, one writes

$$\begin{aligned} I = &\int\limits_{D_y} J_{n-k}(y;\alpha,2k+\gamma+\beta+1,(-1)^k\delta)\; J_{m-\ell}(y,\alpha,2k+\gamma+\beta+1,(-1)^k\delta) \\ &\times \left[\rho_k(y)\rho_\ell(y)|y|^{\beta+\gamma}(1+y)(1-y^2)^{\frac{\alpha-1}{2}}\right] \mathrm{d}y \\ &\qquad \times \int\limits_{D_x} J_k\left(\frac{x}{y};\gamma,\beta,\frac{\delta}{y}\right) J_\ell\left(\frac{x}{y};\gamma,\beta,\frac{\delta}{y}\right) \left[\theta\left(\frac{x}{y}\right)\left(1+\frac{x}{y}\right)\left(\frac{x-\delta}{y}\right)\left(1-\frac{x^2}{y^2}\right)^{\frac{\gamma-1}{2}}\left(\frac{x^2-\delta^2}{y^2}\right)^{\frac{\beta-1}{2}}\right] \mathrm{d}x. \end{aligned}$$

The integral over D_x is directly evaluated using the change of variables $u = x/y$ and comparing with the orthogonality relation (4). The result is thus

$$\begin{aligned} I = h_k(\gamma,\beta)\,\delta_{k\ell} \times &\int\limits_{D_y} J_{n-k}(y;\alpha,2k+\gamma+\beta+1,(-1)^k\delta)\; J_{m-\ell}(y,\alpha,2k+\gamma+\beta+1,(-1)^k\delta) \\ &\qquad \times \rho_k(y)\,\rho_\ell(y) \left[\theta(y)\,(1+y)\,(1-y^2)^{\frac{\alpha-1}{2}}\,\frac{(y^2-\delta^2)^{\frac{2+\gamma+\beta}{2}}}{(y+\delta)}\right] \mathrm{d}y. \end{aligned}$$

Assuming that $k = \ell$ is an even integer, the integral takes the form

$$\begin{aligned} I = h_k(\gamma,\beta)\delta_{k\ell} \times &\int\limits_{D_y} J_{n-k}(y;\alpha,2k+\gamma+\beta+1,\delta)\; J_{m-\ell}(y,\alpha,2k+\gamma+\beta+1,\delta) \\ &\qquad \times (y^2-\delta^2)^k \left[\theta(y)\,(1+y)\,(1-y^2)^{\frac{\alpha-1}{2}}\,\frac{(y^2-\delta^2)^{\frac{2+\gamma+\beta}{2}}}{(y+\delta)}\right] \mathrm{d}y, \end{aligned}$$

which in view of (4) yields

$$I = h_k(\gamma,\beta) h_{n-k}(\alpha,2k+\gamma+\beta+1) \left[\frac{(1-\delta^2)^{\frac{2k+\alpha+\beta+\gamma+3}{2}}}{1+\delta}\right] \delta_{k\ell}\delta_{mn}.$$

Assuming that $k = \ell$ is an odd integer, the integral takes the form

$$\begin{aligned} I = h_k(\gamma,\beta)\delta_{k\ell} \times &\int\limits_{D_y} J_{n-k}(y;\alpha,2k+\gamma+\beta+1,-\delta)\; J_{m-\ell}(y,\alpha,2k+\gamma+\beta+1,-\delta) \\ &\qquad \times (y^2-\delta^2)^{k-1}(y+\delta)^2 \left[\theta(y)\,(1+y)\,(1-y^2)^{\frac{\alpha-1}{2}}\,\frac{(y^2-\delta^2)^{\frac{2+\gamma+\beta}{2}}}{(y+\delta)}\right] \mathrm{d}y, \end{aligned}$$

which given (4) gives

$$I = h_k(\gamma,\beta) h_{n-k}(\alpha,2k+\gamma+\beta+1) \left[\frac{(1-\delta^2)^{\frac{2k+\alpha+\beta+\gamma+3}{2}}}{1-\delta}\right] \delta_{k\ell}\delta_{mn}.$$

Upon combining the k even and k odd cases, one finds (10). This completes the proof. □

It is not difficult to see that the region (12) corresponds to the disjoint union of four triangular domains. The $|\delta| = 1/5$ case is illustrated in Figure 1.



Figure 1: Orthogonality region for $|\delta| = 1/5$

For $\alpha, \beta, \gamma > -1$ and $|\delta| < 1$, it can be verified that the weight function (11) is positive on (12). The orthogonality relation for $|\delta| > 1$ can be obtained in a similar fashion. The result is as follows.

Proposition 2.2. *Let $\alpha, \beta, \gamma > -1$ and $|\delta| > 1$. The two-variable Big -1 Jacobi polynomials defined by* (1) *satisfy the orthogonality relation*

$$\int_{\widetilde{D}_y} \int_{\widetilde{D}_x} \mathscr{J}_{n,k}(x,y)\, \mathscr{J}_{m,\ell}(x,y)\, \widetilde{W}(x,y)\, \mathrm{d}x\, \mathrm{d}y = \widetilde{H}_{nk}\, \delta_{k\ell}\delta_{nm},$$

with respect to the weight function

$$\widetilde{W}(x,y) = \theta(\delta\, x\, y)|y|^{\gamma+\beta}(1+y)\left(1+\frac{x}{y}\right)\left(\frac{\delta - x}{y}\right)(y^2-1)^{\frac{\alpha-1}{2}}\left(\frac{x^2}{y^2}-1\right)^{\frac{\gamma-1}{2}}\left(\frac{\delta^2-x^2}{y^2}\right)^{\frac{\beta-1}{2}}. \tag{13}$$

The integration domain is

$$\widetilde{D}_x = [-|\delta|, -|y|] \cup [|y|, |\delta|], \qquad \widetilde{D}_y = [-|\delta|, 1] \cup [1, |\delta|], \tag{14}$$

and the normalization factor is of the form

$$\widetilde{H}_{nk} = (-1)^k \theta(\delta) \left[\frac{(\delta^2-1)^{\frac{2k+\alpha+\beta+\gamma+3}{2}}}{1+(-1)^k\delta}\right] \widetilde{h}_k(\gamma,\beta)\, \widetilde{h}_{n-k}(\alpha, 2k+\beta+\gamma+1),$$

where $\widetilde{h}_n(a,b)$ is given by (9).

Proof. Similar to proposition 2.1 using instead (7), (8) and (9). □

It can again be seen that the weight function (13) is positive on the domain (14) provided that $\alpha, \beta, \gamma > -1$ and $|\delta| > 1$. The orthogonality region defined by (14) again corresponds to the disjoint union of four triangular domains, as illustrated by Figure 2 for the case $|\delta| = 3$.



Figure 2: Orthogonality region for $|\delta| = 3$

2.1 A special case: the bivariate Little -1 Jacobi polynomials

When $c = 0$, the Big -1 Jacobi polynomials $J_n(x;a,b,c)$ defined by (2) reduce to the so-called Little -1 Jacobi polynomials $j_n(x;a,b)$ introduced in [19]. These polynomials have the hypergeometric representation

$$j_n(x;a,b) = \begin{cases} {}_2F_1\left(\begin{matrix}-\frac{n}{2},\frac{n+a+b+2}{2}\\ \frac{a+1}{2}\end{matrix}\,\middle|\,1-x^2\right) + \frac{n(1-x)}{(a+1)}\,{}_2F_1\left(\begin{matrix}1-\frac{n}{2},\frac{n+a+b+2}{2}\\ \frac{a+3}{2}\end{matrix}\,\middle|\,1-x^2\right), & n \text{ even},\\ {}_2F_1\left(\begin{matrix}-\frac{n-1}{2},\frac{n+a+b+1}{2}\\ \frac{a+1}{2}\end{matrix}\,\middle|\,1-x^2\right) - \frac{(n+a+b+1)(1-x)}{(a+1)}\,{}_2F_1\left(\begin{matrix}-\frac{n-1}{2},\frac{n+a+b+3}{2}\\ \frac{a+3}{2}\end{matrix}\,\middle|\,1-x^2\right), & n \text{ odd}. \end{cases} \tag{15}$$

Taking $\delta = 0$ in (1) leads to the following definition for the two-variable Little -1 Jacobi polynomials:

$$q_{n,k}(x,y) = j_{n-k}(y;\alpha,2k+\beta+\gamma+1)\,y^k\,j_k\left(\frac{x}{y};\gamma,\beta\right), \quad k = 0,1,2\ldots, \quad n = k,k+1,\ldots. \tag{16}$$

It is seen from (16) that the two-variable Little -1 Jacobi polynomials have the structure corresponding to one of the methods to construct bivariate orthogonal polynomials systems proposed by Koornwinder in [13]. For the polynomials (16), the weight function, which can be obtained by taking $\delta = 0$ in either (11) or (13), can also be recovered using the general scheme given in [13]. For $\delta = 0$, the region (12) reduces to two vertically opposite triangles.

3 Bispectrality of the bivariate Big -1 Jacobi polynomials

In this section, the two-variable Big -1 Jacobi polynomials $\mathscr{J}_{n,k}(x,y)$ are shown be the joint eigenfunctions of a pair of first-order differential operators involving reflections. Their recurrence relations are also derived. The results are obtained through a limiting process from the corresponding properties of the two-variable Big q-Jacobi polynomials introduced by Lewanowicz and Woźny [16].

3.1 Bivariate Big q-Jacobi polynomials

Let us review some of the properties of the bivariate q-polynomials introduced in [16]. The two-variable Big q-Jacobi polynomials, denoted $\mathscr{P}_{n,k}(x,y;a,b,c,d;q)$ are defined as

$$\mathscr{P}_{n,k}(x,y;a,b,c,d;q) = P_{n-k}(y;a,bcq^{2k+1},dq^k;q)\,y^k\,\left(\frac{dq}{y};q\right)_k\,P_k\left(\frac{x}{y};c,b,\frac{d}{y};q\right), \tag{17}$$

where $(a;q)_n$ stands for the q-Pochhammer symbol [4] and where $P_n(x;a,b,c;q)$ are the Big q-Jacobi polynomials [12]. The two-variable Big q-Jacobi polynomials satisfy the eigenvalue equation [16]

$$\Omega \mathscr{P}_{n,k}(x,y) = \left[\frac{q^{1-n}(q^n-1)(abcq^{n+2}-1)}{(q-1)^2}\right]\mathscr{P}_{n,k}(x,y), \tag{18}$$

where Ω is the q-difference operator

$$\begin{aligned}\Omega &= (x-dq)(x-acq^2)\mathbf{D}_{q,x}\mathbf{D}_{q^{-1},x} + (y-aq)(y-dq)\mathbf{D}_{q,y}\mathbf{D}_{q^{-1},y}\\ &+ q^{-1}(x-dq)(y-aq)\mathbf{D}_{q^{-1},x}\mathbf{D}_{q^{-1},y} + acq^3(bx-d)(y-1)\mathbf{D}_{q,x}\mathbf{D}_{q,y}\\ &+ \frac{(abcq^3-1)(x-1)-(acq^2-1)(dq-1)}{q-1}\mathbf{D}_{q,x} + \frac{(abcq^3-1)(y-1)-(aq-1)(dq-1)}{q-1}\mathbf{D}_{q,y},\end{aligned}$$

and where $\mathbf{D}_{q,x}$ stands for the q-derivative

$$\mathbf{D}_{q,x}f(x,y) = \frac{f(qx,y)-f(x,y)}{x(q-1)}.$$

The bivariate Big q-Jacobi polynomials also satisfy the pair of recurrence relations [16]

$$\begin{aligned}y\mathscr{P}_{n,k}(x,y) &= a_{nk}\mathscr{P}_{n+1,k}(x,y) + b_{nk}\mathscr{P}_{n,k}(x,y) + c_{nk}\mathscr{P}_{n-1,k}(x,y),\\ x\mathscr{P}_{n,k}(x,y) &= e_{nk}\mathscr{P}_{n+1,k-1}(x,y) + f_{nk}\mathscr{P}_{n+1,k}(x,y) + g_{nk}\mathscr{P}_{n+1,k+1}(x,y)\\ &+ r_{nk}\mathscr{P}_{n,k-1}(x,y) + s_{nk}\mathscr{P}_{n,k}(x,y) + t_{nk}\mathscr{P}_{n,k+1}(x,y)\\ &+ u_{nk}\mathscr{P}_{n-1,k-1}(x,y) + v_{nk}\mathscr{P}_{n-1,k}(x,y) + w_{nk}\mathscr{P}_{n-1,k+1}(x,y),\end{aligned} \tag{19}$$

where the recurrence coefficients read

$$\begin{aligned}a_{nk} &= \frac{(1-aq^{n-k+1})(1-abcq^{n+k+2})(1-dq^{n+1})}{(abcq^{2n+2};q)_2}, \quad & b_{nk} &= 1-a_{nk}-c_{nk},\\ w_{nk} &= \frac{abc\sigma_k q^{n+2k+3}(abcq^{n+1}-d)(q^{n-k-1};q)_2}{(1-dq^{k+1})(abcq^{2n+1};q)_2}, \quad & f_{nk} &= a_{nk}(bcq^k\tau_k-\sigma_k+1),\\ e_{nk} &= \frac{\tau_k bcq^k(dq^k-1)(1-dq^{n+1})(aq^{n-k+1};q)_2}{(abcq^{2n+2};q)_2}, \quad & v_{nk} &= c_{nk}(bcq^k\tau_k-\sigma_k+1),\\ g_{nk} &= \frac{\sigma_k(1-dq^{n+1})(abcq^{n+k+2};q)_2}{(1-dq^{k+1})(abcq^{2n+2};q)_2}, & s_{nk} &= b_{nk}(bcq^k\tau_k-\sigma_k+1)+d(q^{k+1}\sigma_k-\tau_k),\end{aligned}$$

with

$$\begin{aligned}t_{nk} &= \frac{q^{k+1}\sigma_k z_n(1-q^{n-k})(1-abcq^{n+k+2})}{1-dq^{k+1}},\\ r_{nk} &= \tau_k z_n(dq^k-1)(1-aq^{n-k+1})(1-bcq^{n+k+1}),\\ u_{nk} &= \frac{\tau_k aq^{n-k+1}(dq^k-1)(abcq^{n+1}-d)(bcq^{n+k};q)_2}{(abcq^{2n+1};q)_2},\\ c_{nk} &= \frac{adq^{n+1}(q^{n-k}-1)(1-bcq^{n+k+1})(1-abcd^{-1}q^{n+1})}{(abcq^{2n+1};q)_2},\end{aligned}$$

and where σ_k, τ_k and z_n are given by

$$\begin{aligned}\sigma_k &= \frac{(1-cq^{k+1})(1-bcq^{k+1})}{(bcq^{2k+1};q)_2}, \quad & \tau_k &= -\frac{cq^{k+1}(1-q^k)(1-bq^k)}{(bcq^{2k};q)_2},\\ z_n &= \frac{abcq^{n+1}(1+q-dq^{n+1})-d}{(1-abcq^{2n+1})(1-abcq^{2n+3})}.\end{aligned}$$

3.2 $\mathscr{J}_{n,k}(x,y)$ as a $q \to -1$ limit of $\mathscr{P}_{n,k}(x,y)$

The two-variable Big -1 Jacobi polynomials $\mathscr{J}_{n,k}(x,y)$ can be obtained from the bivariate Big q-Jacobi by taking $q \to -1$. Indeed, a direct calculation using the expression (17) shows that

$$\lim_{\epsilon \to 0} \mathscr{P}_{n,k}(x,y;-e^{\epsilon\alpha},-e^{\epsilon\beta},-e^{\epsilon\gamma},\delta,-e^{\epsilon}) = \mathscr{J}_{n,k}(x,y;\alpha,\beta,\gamma,\delta), \tag{20}$$

where we have used the notation $\mathscr{J}_{n,k}(x,y;\alpha,\beta,\gamma,\delta)$ to exhibit the parameters appearing in the Big -1 Jacobi polynomials defined in (1). A similar limit was considered in [20] to obtain the univariate Big -1 Jacobi polynomials in terms of the Big q-Jacobi polynomials.

3.3 Eigenvalue equation for the Big -1 Jacobi polynomials

The eigenvalue equation (18) for the Big q-Jacobi polynomials and the relation (20) between the Big q-Jacobi polynomials and the Big -1 Jacobi polynomials can be used to obtain an eigenvalue equation for the latter.

Proposition 3.1. *Let L_1 be the first-order differential / difference operator*

$$\begin{aligned} L_1 = {} & G_5(x,y)R_xR_y\partial_y + G_6(x,y)R_y\partial_y + G_7(x,y)R_xR_y\partial_x + G_8(x,y)R_x\partial_x \\ & + G_1(x,y)R_xR_y + G_2(x,y)R_x + G_3(x,y)R_y - (G_1(x,y)+G_2(x,y)+G_3(x,y))\mathbb{I}, \end{aligned} \tag{21}$$

where R_x, R_y are reflection operators and where the coefficients read

$$G_1(x,y) = \frac{x[1+\beta+\gamma-y(\alpha+\beta+\gamma+2)]-\delta[y(\alpha+\gamma+1)-\gamma]}{4xy}, \quad G_8(x,y) = \frac{(\delta+x)(x-y)}{2xy}, \tag{22a}$$

$$G_2(x,y) = -\frac{x[x(\beta+\gamma+1)-\beta y]+\delta(y+\gamma x)}{4x^2y}, \qquad G_7(x,y) = \frac{(\delta+x)(y-1)}{2y}, \tag{22b}$$

$$G_3(x,y) = -\delta\left(\frac{x+\alpha xy-y[y(\alpha+\gamma+1)-\gamma]}{4xy^2}\right) \qquad G_6(x,y) = \frac{\delta(x-y)(y-1)}{2xy}, \tag{22c}$$

$$G_5(x,y) = \frac{(\delta+x)(y-1)}{2x}. \tag{22d}$$

The Big -1 Jacobi polynomials satisfy the eigenvalue equation

$$L_1\mathscr{P}_{n,k}(x,y) = \mu_n\mathscr{P}_{n,k}(x,y), \qquad \mu_n = \begin{cases} -\frac{n}{2}, & n \text{ even}, \\ \frac{n+\alpha+\beta+\gamma+2}{2}, & n \text{ odd}. \end{cases}$$

Furthermore, let L_2 be the differential / difference operator

$$L_2 = \frac{2(y-x)(x+\delta)}{x}R_x\partial_x + \frac{(\gamma+\beta+1)x^2+(\delta\gamma-\beta y)x+\delta y}{x^2}(R_x-\mathbb{I}),$$

The Big -1 Jacobi polynomials satisfy the eigenvalue equation

$$L_2\mathscr{P}_{n,k}(x,y) = \nu_k\mathscr{P}_{n,k}(x,y), \qquad \nu_k = \begin{cases} 2k, & k \text{ even}, \\ -2(k+\beta+\gamma+1), & k \text{ odd}, \end{cases}.$$

Proof. The eigenvalue equation with respect to L_1 is obtained by dividing both sides of (18) by $(1+q)$ and taking the $q \to -1$ limit according to (20). The eigenvalue equation with respect to L_2 is obtained by combining (1), (2) and (3). □

The two-variable Big -1 Jacobi polynomials are thus the joint eigenfunctions of the first order differential operators with reflections L_1 and L_2. It is directly verified that these operators commute with one another, as should be.

The $q \to -1$ limit (20) of the recurrence relations (19) can also be taken to obtain the recurrence relations satisfied by the Big -1 Jacobi polynomials. The result is as follows.

Proposition 3.2. *The Big -1 Jacobi polynomials satisfy the recurrence relations*

$$
\begin{aligned}
&y\,\mathscr{J}_{n,k}(x,y) = \widetilde{a}_{nk}\,\mathscr{J}_{n+1,k}(x,y) + \widetilde{b}_{nk}\,\mathscr{J}_{n,k}(x,y) + \widetilde{c}_{nk}\,\mathscr{J}_{n-1,k}(x,y),\\
&x\,\mathscr{J}_{n,k}(x,y) = \widetilde{e}_{nk}\,\mathscr{J}_{n+1,k-1}(x,y) + \widetilde{f}_{nk}\,\mathscr{J}_{n+1,k}(x,y) + \widetilde{g}_{nk}\,\mathscr{J}_{n+1,k+1}(x,y)\\
&\quad + \widetilde{r}_{nk}\,\mathscr{J}_{n,k-1}(x,y) + \widetilde{s}_{nk}\,\mathscr{J}_{n,k}(x,y) + \widetilde{t}_{nk}\,\mathscr{J}_{n,k+1}(x,y)\\
&\qquad\qquad + \widetilde{u}_{nk}\,\mathscr{J}_{n-1,k-1}(x,y) + \widetilde{v}_{nk}\,\mathscr{J}_{n-1,k}(x,y) + \widetilde{w}_{nk}\,\mathscr{J}_{n-1,k+1}(x,y).
\end{aligned}
$$

With

$$
\widetilde{\tau}_k = \frac{k+\beta\phi_k}{2k+\beta+\gamma}, \quad \widetilde{\sigma}_k = \frac{k+\beta\phi_k+\gamma+1}{2k+\beta+\gamma+2}, \quad \widetilde{z}_n = \frac{(-1)^n - \delta(2n+\alpha+\beta+\gamma+2)}{(2n+\alpha+\beta+\gamma+1)(2n+\alpha+\beta+\gamma+3)},
$$

where $\phi_k = (1-(-1)^k)/2$ is the characteristic function for odd numbers, the recurrence coefficients read

$$
\widetilde{a}_{n,k} = \frac{1+\delta_n}{2n+\alpha+\beta+\gamma+3} \times \begin{cases} n-k+\alpha+1, & n+k \text{ even},\\ n+k+\alpha+\beta+\gamma+2, & n+k \text{ odd}, \end{cases}
$$

$$
\widetilde{c}_{n,k} = \frac{1+\delta_{n+1}}{2n+\alpha+\beta+\gamma+1} \times \begin{cases} n-k, & n+k \text{ even},\\ n+k+\beta+\gamma+1, & n+k \text{ odd}, \end{cases}
$$

$$
\widetilde{e}_{n,k} = \frac{\widetilde{\tau}_k(1-\delta_k)(1+\delta_n)}{2n+\alpha+\beta+\gamma+3} \times \begin{cases} n-k+\alpha+1, & n+k \text{ even},\\ n-k+\alpha+2, & n+k \text{ odd}, \end{cases}
$$

$$
\widetilde{g}_{n,k} = \frac{\widetilde{\sigma}_k(1+\delta_n)}{(1+\delta_k)(2n+\alpha+\beta+\gamma+3)} \times \begin{cases} n+k+\alpha+\beta+\gamma+3, & n+k \text{ even},\\ n+k+\alpha+\beta+\gamma+2, & n+k \text{ odd}, \end{cases}
$$

$$
\widetilde{r}_{n,k} = 2\widetilde{\tau}_k\widetilde{z}_n((-1)^k-\delta) \times \begin{cases} n-k+\alpha+1 & n+k \text{ even}\\ n+k+\beta+\gamma+1 & n+k \text{ odd} \end{cases}
$$

$$
\widetilde{t}_{n,k} = \frac{2(-1)^{k+1}\widetilde{\sigma}_k\widetilde{z}_n}{1+\delta_k} \times \begin{cases} n-k, & n+k \text{ even},\\ n+k+\alpha+\beta+\gamma+2, & n+k \text{ odd}, \end{cases}
$$

$$
\widetilde{u}_{nk} = \frac{\widetilde{\tau}_k(1-\delta_k)(1-\delta_n)}{(2n+\alpha+\beta+\gamma+1)} \times \begin{cases} n+k+\beta+\gamma, & n+k \text{ even},\\ n+k+\beta+\gamma+1, & n+k \text{ odd}, \end{cases}
$$

$$
\widetilde{w}_{n,k} = \frac{\widetilde{\sigma}_k(1-\delta_n)}{(1+\delta_k)(2n+\alpha+\beta+\gamma+1)} \times \begin{cases} n-k, & n+k \text{ even},\\ n-k-1, & n+k \text{ odd}, \end{cases}
$$

with $\delta_n = (-1)^n\delta$ and

$$
\begin{aligned}
&\widetilde{b}_{n,k} = 1-\widetilde{a}_{n,k}-\widetilde{c}_{n,k}, \quad \widetilde{f}_{n,k} = \widetilde{a}_{n,k}(1-\widetilde{\sigma}_k-\widetilde{\tau}_{n,k}), \quad \widetilde{s}_{n,k} = \widetilde{b}_{n,k}(1-\widetilde{\sigma}_k-\widetilde{\tau}_k) - \delta_k(\widetilde{\sigma}_k-\widetilde{\tau}_k)\\
&\widetilde{v}_{n,k} = \widetilde{c}_{n,k}(1-\widetilde{\sigma}_k-\widetilde{\tau}_k)
\end{aligned}
$$

Proof. The result is obtained by applying the limit (20) to the recurrence relations (19). □

4 A Pearson-type system for Big -1 Jacobi

Let us now show how the weight function $W(x,y)$ for the polynomials $\mathscr{J}_{n,k}(x,y)$ can be recovered as the symmetry factor for the operator L_1 given in (21). The symmetrization condition for L_1 is

$$(W(x,y)L_1)^* = W(x,y)L_1, \tag{23}$$

where M^* denotes the Lagrange adjoint. For an operator of the form

$$M = \sum_{\mu,\nu,k,j} A_{k,j}(x,y)\,\partial_x^k\,\partial_y^j\,R_x^\mu\,R_y^\nu,$$

for $\mu,\nu \in \{0,1\}$ and $k,j = 0,1,2,\ldots$, the Lagrange adjoint reads

$$M^* = \sum_{\mu,\nu,k,j} (-1)^{k+j}\,R_x^\mu R_y^\nu\,\partial_y^j\,\partial_x^k\,A_{k,j}(x,y),$$

where we have assumed that $W(x,y)$ is defined on a symmetric region with respect to R_x and R_y. Imposing the condition (23), one finds the following system of Pearson-type equations:

$$W(x,y)G_8(x,y) = W(-x,y)G_8(-x,y), \tag{24a}$$
$$W(x,y)G_7(x,y) = W(-x,-y)G_7(-x,-y), \tag{24b}$$
$$W(x,y)G_6(x,y) = W(x,-y)G_6(x,-y), \tag{24c}$$
$$W(x,y)G_5(x,y) = W(-x,-y)G_5(-x,-y) \tag{24d}$$
$$W(x,y)G_3(x,y) = W(x,-y)G_3(x,-y) - \partial_y(W(x,-y)G_6(x,-y)), \tag{24e}$$
$$W(x,y)G_2(x,y) = W(-x,y)G_2(-x,y) - \partial_x(W(-x,y)G_8(-x,y)), \tag{24f}$$
$$\begin{aligned} W(x,y)G_1(x,y) = {} & W(-x,-y)G_1(-x,-y) \\ & - \partial_x(W(-x,-y)G_7(-x,-y)) - \partial_y(W(-x,-y)G_5(-x,-y)), \end{aligned} \tag{24g}$$

where the functions $G_i(x,y)$, $i = 1,\ldots,8$, are given by (22). We assume that $W(x,y) > 0$ and moreover that $|\delta| \leqslant |x| \leqslant y \leqslant 1$. Upon substituting (22) in (24a), one finds

$$(x+\delta)(x-y)W(x,y) = -(x-\delta)(x+y)W(-x,y),$$

for which the general solution is of the form

$$W(x,y) = \theta(x)(x-\delta)(x+y)\,f_1(x^2,y), \tag{25}$$

where f_1 is an arbitrary function. Using (25) in (24b) yields

$$(y-1)\,f_1(x^2,y) = (y+1)\,f_1(x^2,-y),$$

which has the general solution

$$f_1(x^2,y) = \theta(y)(y+1)\,f_2(x^2,y^2),$$

where f_2 is an arbitrary function. The function $W(x,y)$ is thus of the general form

$$W(x,y) = \theta(x)\theta(y)(x-\delta)(x+y)(y+1)\,f_2(x^2,y^2). \tag{26}$$

Upon substituting the above expression in (24f), one finds after simplifications

$$\left[\frac{\beta-1}{x^2-\delta^2} + \frac{\gamma-1}{x^2-y^2}\right] x\,f_2(x^2,y^2) = \partial_x\,f_2(x^2,y^2).$$

After separation of variable, the result is

$$f_2(x^2,y^2) = (x^2-\delta^2)^{\frac{\beta-1}{2}}(y^2-x^2)^{\frac{\gamma-1}{2}} f_3(y^2).$$

Finally, upon substituting the above equation in (24e) one finds

$$y(\alpha-1)f_3(y^2) = (y^2-1)\partial_y f_3(y^2),$$

which gives $f_3 = (1-y^2)^{\frac{\alpha-1}{2}}$ and thus

$$f_2(x^2,y^2) = (x^2-\delta^2)^{\frac{\beta-1}{2}}(y^2-x^2)^{\frac{\gamma-1}{2}}(1-y^2)^{\frac{\alpha-1}{2}}.$$

Upon combining the above expression with (26), we find

$$W(x,y) = \theta(xy)|y|^{\beta+\gamma}(1+y)\left(1+\frac{x}{y}\right)\left(\frac{x-\delta}{y}\right)(1-y^2)^{\frac{\alpha-1}{2}}\left(1-\frac{x^2}{y^2}\right)^{\frac{\gamma-1}{2}}\left(\frac{x^2-\delta^2}{y^2}\right)^{\frac{\beta-1}{2}},$$

which indeed corresponds to the weight function of proposition 2.1. The weight function $W(x,y)$ for the two-variable Big -1 Jacobi polynomials thus corresponds to the symmetry factor for L_1.

5 Conclusion

In this paper, we have introduced and characterized a new family of two-variable orthogonal polynomials that generalize of the Big -1 Jacobi polynomials. We have constructed their orthogonality measure and we have derived explicitly their bispectral properties. We have furthermore shown that the weight function for these two-variable polynomials can be recovered by symmetrization of the first-order differential operator with reflections that these polynomials diagonalize. The two-variable orthogonal polynomials introduced here are the first example of the multivariate generalization of the Bannai-Ito scheme. It would be of great interest to construct multivariate extensions of the other families of polynomials of this scheme.

Acknowledgments

VXG holds an Alexander-Graham-Bell fellowship from the Natural Science and Engineering Research Council of Canada (NSERC). JML holds a scholarship from NSERC. The research of LV is supported in part by NSERC. AZ would like to thank the Centre de recherches mathématiques for its hospitality.

A Normalization coefficients for Big -1 Jacobi polynomials

In this appendix, we present a derivation of the normalization coefficients (6) and (9) appearing in the orthogonality relation of the univariate Big -1 Jacobi polynomials. The result is obtained by using their kernel partners, the Chihara polynomials. These polynomials, denoted by $C_n(x;\alpha,\beta,\gamma)$, have the expression [8]

$$C_{2n}\left(x;\alpha,\beta,\gamma\right) = (-1)^n\frac{(\alpha+1)_n}{(n+\alpha+\beta+1)_n}\,{}_2F_1\left(\begin{matrix}-n, n+\alpha+\beta+1\\ \alpha+1\end{matrix}\,\middle|\, x^2-\gamma^2\right),$$
$$C_{2n+1}\left(x;\alpha,\beta,\gamma\right) = (-1)^n\frac{(\alpha+2)_n}{(n+\alpha+\beta+2)_n}(x-\gamma)\,{}_2F_1\left(\begin{matrix}-n, n+\alpha+\beta+2\\ \alpha+2\end{matrix}\,\middle|\, x^2-\gamma^2\right).$$

For $\alpha, \beta > -1$, they satisfy the orthogonality relation

$$\int_{\mathscr{E}} C_n\left(x;\alpha,\beta,\gamma\right) C_m\left(x;\alpha,\beta,\gamma\right) \theta(x)(x+\gamma)(x^2-\gamma^2)^{\alpha}(1+\gamma^2-x^2)^{\beta}\,\mathrm{d}x = \eta_n \delta_{nm}, \tag{27}$$

on the interval $\mathscr{E} = [-\sqrt{1+\gamma^2}, -|\gamma|] \cup [|\gamma|, \sqrt{1+\gamma^2}]$ and their normalization coefficients read

$$\eta_{2n} = \frac{\Gamma(n+\alpha+1)\Gamma(n+\beta+1)}{\Gamma(n+\alpha+\beta+1)} \frac{n!}{(2n+\alpha+\beta+1)[(n+\alpha+\beta+1)_n]^2},$$
$$\eta_{2n+1} = \frac{\Gamma(n+\alpha+2)\Gamma(n+\beta+1)}{\Gamma(n+\alpha+\beta+2)} \frac{n!}{(2n+\alpha+\beta+2)[(n+\alpha+\beta+2)_n]^2}.$$

Let $\widehat{J}_n(x)$ be the monic Big -1 Jacobi polynomials:

$$\widehat{J}_n(x) = \kappa_n J_n(x) = x^n + \mathscr{O}(x^{n-1}),$$

where κ_n is given by

$$\kappa_n(a,b,c) = \begin{cases} \frac{(1-c^2)^{\frac{n}{2}} \left(\frac{a+1}{2}\right)_{\frac{n}{2}}}{\left(\frac{n+a+b+2}{2}\right)_{\frac{n}{2}}}, & n \text{ even}, \\ \frac{(1+c)(1-c^2)^{\frac{n-1}{2}} \left(\frac{a+1}{2}\right)_{\frac{n+1}{2}}}{\left(\frac{n+a+b+1}{2}\right)_{\frac{n+1}{2}}}, & n \text{ odd}. \end{cases}$$

The kernel polynomials $\widetilde{K}_n(x;a,b,c)$ associated to $\widehat{J}_n(x;a,b,c)$ are defined through the Christoffel transformation [2]:

$$\widetilde{K}_n(x;a,b,c) = \frac{\widehat{J}_{n+1}(x) - \frac{\widehat{J}_{n+1}(\nu)}{\widehat{J}_n(\nu)} \widehat{J}_n(x)}{x-\nu}. \tag{28}$$

For $\nu = 1$, the monic polynomials $\widetilde{K}_n(x;a,b,c)$ can be expressed in terms of the Chihara polynomials. Indeed, one can show that [8]

$$\widetilde{K}_n(x;a,b,c) = (\sqrt{1-c^2})^n\, C_n\left(\frac{x}{\sqrt{1-c^2}}; \frac{b-1}{2}, \frac{a+1}{2}, \frac{-c}{\sqrt{1-c^2}}\right). \tag{29}$$

Let $\mathscr{M}$ be the orthogonality functional for the Big -1 Jacobi polynomials. In view of the relation (28), one can write [2]

$$\mathscr{M}[(x-\nu)\,\widetilde{K}_n(x;a,b,c)\,x^k] = -\frac{\widehat{J}_{n+1}(\nu)}{\widehat{J}_n(\nu)} \mathscr{M}[\widehat{J}_n^2(x;a,b,c)]\delta_{kn}.$$

By linearity, one thus finds

$$\widetilde{\eta}_n = -\frac{\widehat{J}_{n+1}(\nu)}{\widehat{J}_n(\nu)} \hat{h}_n.$$

where $\widetilde{\eta}_n = \mathscr{M}[(x-\nu)\,\widetilde{K}_n^2(x)]$ and $\hat{h}_n = \mathscr{M}[\widehat{P}_n^2(x)]$. For $\nu = 1$, the value of $\widetilde{\eta}_n$ is easily computed from (27) and (29). The above relation then gives the desired coefficients.

References

[1] G. Andrews, R. Askey, and R. Roy. *Special functions*. Cambridge University Press, 1999.

[2] T. Chihara. *An Introduction to Orthogonal Polynomials*. Dover Publications, 2011.

[3] C. Dunkl. Orthogonal polynomials in two variables of q-Hahn and q-Jacobi type. *SIAM J. Algebr. Discrete Methods*, 1:137–151, 1980.

[4] G. Gasper and M. Rahman. *Basic Hypergeometric Series*. Encyclopedia of Mathematics and its Applications. Cambridge University Press, 2^{nd} edition, 2004.

[5] V. X. Genest, M. Ismail, L. Vinet, and A. Zhedanov. The Dunkl oscillator in the plane: I. Superintegrability, separated wavefunctions and overlap coefficients. *J. Phys. A: Math. Theor.*, 46:145201, 2013.

[6] V. X. Genest, M. Ismail, L. Vinet, and A. Zhedanov. The Dunkl oscillator in the plane II: Representations of the symmetry algebra. *Comm. Math. Phys.*, 329:999–1029, 2014.

[7] V. X. Genest, L. Vinet, and A. Zhedanov. The singular and the $2:1$ anisotropic Dunkl oscillators in the plane. *J. Phys. A: Math. Theor.*, 46:325201, 2013.

[8] V. X. Genest, L. Vinet, and A. Zhedanov. A "continuous" limit of the Complementary Bannai-Ito polynomials: Chihara polynomials. *SIGMA*, 10:38–55, 2014.

[9] V. X. Genest, L. Vinet, and A. Zhedanov. A Laplace-Dunkl equation on S^2 and the Bannai-Ito algebra. *Comm. Math. Phys. (to appear)*, 2015.

[10] V.X. Genest, L. Vinet, and A. Zhedanov. Bispectrality of the Complementary Bannai–Ito polynomials. *SIGMA*, 9:18–37, 2013.

[11] J. Harnad, L. Vinet, O. Yermolayeva, and A. Zhedanov. Two-dimensional Krall-Sheffer polynomials and integrable systems. *J. Phys. A: Math. Gen.*, 34:10619–10625, 2001.

[12] R. Koekoek, P.A. Lesky, and R.F. Swarttouw. *Hypergeometric orthogonal polynomials and their q-analogues*. Springer, 1^{st} edition, 2010.

[13] T. Koornwinder. Two-variable analogues of the classical orthogonal polynomials. In R. Askey, editor, *Theory and applications of special functions*, pages 435–495. Academic Press, 1975.

[14] H. L. Krall and I. M. Sheffer. Orthogonal polynomials in two variables. *Ann. Mat. Pura Appl.*, 76:325–376, 1967.

[15] K. H. Kwon, J. K. Lee, and L. L. Littlejohn. Orthogonal polynomial eigenfunctions of second-order partial differential equations. *Trans. Amer. Math. Soc.*, 353:3629–3647, 2001.

[16] S. Lewanowicz and P. Woźny. Two-variable orthogonal polynomials of q-Jacobi type. *J. Comput. Appl. Math.*, 233, 2010.

[17] W. Miller, S. Post, and P. Winternitz. Classical and quantum superintegrability with applications. *J. Phys. A: Math. Theor.*, 46:423001, 2013.

[18] S. Tsujimoto, L. Vinet, and A. Zhedanov. Dunkl shift operators and Bannai–Ito polynomials. *Adv. Math.*, 229:2123–2158, 2012.

[19] L. Vinet and A. Zhedanov. A 'missing' family of classical orthogonal polynomials. *J. Phys. A: Math. Theor.*, 44:085201, 2011.

[20] L. Vinet and A. Zhedanov. A limit $q = -1$ for the Big q-Jacobi polynomials. *Trans. Amer. Math. Soc.*, 364:5491–5507, 2012.